

Signal Heading:	<p style="text-align: center;">POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING</p>
------------------------	--

Product Name:	[INSERT HERE]
----------------------	---------------

Constituent Statement:	<p>300 g/L 2,4-D present as the 2-ethylhexyl ester 6.25 g/L FLORASULAM</p> <p>[Applicants if your product contains other scheduled excipients please include these here or delete this text]</p>
-------------------------------	--

Mode of Action:	GROUP B I HERBICIDE
------------------------	----------------------------

Statement of Claims:	<p>[INSERT HERE]</p> <p>THIS IS A PHENOXY HERBICIDE THAT CAN CAUSE SEVERE DAMAGE TO NATIVE VEGETATION AND SUSCEPTIBLE CROPS SUCH AS COTTON, GRAPES, TOMATOES, OILSEED CROPS AND ORNAMENTALS.</p>
-----------------------------	--

Net Contents:	[INSERT HERE]
----------------------	---------------

Restrains:	See 'SPRAY DRIFT RESTRAINTS' and 'GENERAL RESTRAINTS' sections. This can be uploaded as an attachment when submitting a label
-------------------	---

Directions For Use:	<p>Please submit a directions for use table.</p> <p>As your product is the only registered group 20B product, and because none of the uses for your product are expected to be impacted by the review an updated directions for use has not been prepared for your product.</p>
----------------------------	---

Other Limitations:	IN TASMANIA, THIS PRODUCT MAY ONLY BE USED FROM 15 APRIL TO 15 SEPTEMBER UNLESS OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.
---------------------------	---

Withholding Period:	<p>WITHOLDING PERIODS: PASTURE, CEREAL CROPS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION CROP HARVEST WITHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.</p>
----------------------------	---

Trade Advice:	
----------------------	--

General Instructions:	[INSERT HERE]
------------------------------	---------------

Resistance Warning:	Resistant Weeds Warning GROUP I HERBICIDE
----------------------------	--

	<p>[INSERT PRODUCT NAME] is a member of the triazolopyrimidine sulfonanilide and phenoxy groups of herbicides. The product has the disruptors of plant cell growth and acetolactate synthase (ALS) inhibitor modes of action.</p> <p>For weed resistance management [INSERT PRODUCT NAME] is a Group B and I herbicide. Some naturally-occurring weed biotypes resistant to [INSERT PRODUCT NAME] and other Group B and I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by [INSERT PRODUCT NAME] or other Group B and I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, [INSERT COMPANY NAME] accepts no liability for any losses that may result from the failure of [INSERT PRODUCT NAME] to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant or local Department of Agriculture.</p> <p>The herbicide is a broadleaf herbicide with no grass weed activity and exerts no selection pressure on annual ryegrass.</p>
--	--

<p>Precautions:</p>	<p>RE-ENTRY</p> <p>Do not allow entry into treated areas until the spray has dried when conducting low exposure activities such as hand weeding unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.</p> <p>Do not allow entry into treated areas for 13 days when conducting medium exposure activities such as scouting unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.</p>
----------------------------	---

<p>Protection Statements:</p>	<p>PROTECTION OF CROPS, NATIVE AND NON-TARGET PLANTS</p> <p>DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.</p> <p>Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</p> <p>Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.</p> <p>INTEGRATED PEST MANAGEMENT</p> <p>Toxic to beneficial arthropods. Not compatible with integrated pest management (IPM) programs utilising beneficial arthropods. Minimise spray drift to reduce harmful effects on beneficial arthropods in non-crop areas.</p>
--------------------------------------	---

<p>Storage and Disposal:</p>	<p>[Applicants, please chose from the relevant statements below.]</p> <p>STORAGE AND DISPOSAL:</p> <p>Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.</p>
-------------------------------------	---

	<p>[Refillable Containers] Empty contents fully into application equipment. Close all valves and return to [point of supply/designated collection point/other specific collection details] for refill or storage.</p> <p>[Metal drums and plastic containers] Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.</p> <p>If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p> <p>[Containers accompanied by a leaflet If a label leaflet is to be used as part of the label for various containers that have different disposal instructions, the following 'Disposal' statement may be shown on the label leaflet in place of the full disposal instructions:]</p> <p>The method of disposal of the container depends on the container type. Read the 'Storage and disposal' instructions on the label that is attached to the container.</p> <p>[drumMUSTER containers] This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately.</p>
--	---

<p>Safety Directions:</p>	<p>Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with skin</p> <p>When opening the container and preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). In addition, when mixing and loading, wear elbow-length chemical resistant gloves.</p> <p>Wash hands after use. After each day's use wash gloves and contaminated clothing.</p>
----------------------------------	--

<p>First Aid Instructions:</p>	<p>[Applicants if your product contains other scheduled excipients additional first aid instructions may be required, the first aid instructions given here are for the 2,4-D active only]</p> <p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.</p>
---------------------------------------	--

Restraints

GENERAL RESTRAINTS

DO NOT exceed maximum application rate of 15 L/ha (4500 g ae/ha)

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of runoff for at least 3 days after application.

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

[Applicants please INSERT additional restraints not covered by the proposed regulatory decision if they are applicable to your product]

SPRAY DRIFT RESTRAINTS

DO NOT apply by a vertical sprayer.

DO NOT apply by aircraft.

Specific definitions for terms used in this section of the label can be found at www.apvma.gov.au/spraydrift

DO NOT allow **bystanders** to come into contact with the spray cloud

DO NOT apply in a manner that may cause an **unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production**, or cause contamination of plant or livestock commodities, outside the application site from **spray drift**. The **buffer zones** in the relevant buffer zone tables below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas

DO NOT apply unless the **wind speed** is between 3 and 20 kilometres per hour at the **application site** during the time of application.

DO NOT apply if there are **hazardous surface temperature inversion** conditions present at the **application site** during the time of application. **Surface temperature inversion conditions** exist most evenings one to two hours before sunset and persist until one or two hours after sunrise.

BOOM SPRAYERS

DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets not smaller than a VERY COARSE **spray droplet size category**
- minimum distances between the **application site** and downwind **sensitive areas** (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for boom sprayers') are observed

Buffer Zones for Boom Sprayers

Application rate (/ha)		Mandatory buffer zones (distances given in metres)
------------------------	--	--

	Boom Height above target canopy	Bystander Areas	Natural Aquatic Areas	Pollinator Areas	Vegetation Areas	Livestock Areas
Up to 1 L (300 g a/ha)	0.5m or lower	Not Required	Not Required	Not Required	10	Not Required
	1.0m or lower		25		35	

DRAFT